



# An Integrated Multi-Scale Approach to the Study of Evapotranspiration on the Alaskan North Slope: Preliminary Characterization of Fluxes and Turbulence in the Imnavait Creek



Chris Wyatt<sup>1</sup>, John Mumm<sup>2</sup>, Erin Trochim<sup>2</sup>, Javier Fochesatto<sup>1</sup>, Anupma Prakash<sup>1</sup>, Martha Anderson<sup>3</sup>, Doug Kane<sup>2</sup>

(1) Geophysical Institute, University of Alaska Fairbanks. AK 99775. (2) Water and Environmental Research Center, University of Alaska Fairbanks


(3) United States Department of Agriculture

American Geophysical Union, Fall Meeting, San Francisco. December, 2009

Session Understanding Land-Atmosphere Interactions with Models & Observations N°: H41G-0980

## Introduction

Evapotranspiration (ET) plays a significant role in the hydrologic cycle of Arctic basins. Surface-atmosphere exchanges due to ET in the Imnavait Creek Basin are estimated from water balance computations to be about 74% of summer precipitation or 50% of annual precipitation. Even though ET is a significant component of the hydrologic cycle in this region, the bulk estimates don't accurately account for spatial and temporal variability due to vegetation type, topography, etc. (Kane and Yang, 2004). A preliminary experiment was carried out in the summer of 2009 to characterize the turbulent fluxes (i.e. buoyancy fluxes) at two levels of 1 and 3 m AGL and the heat fluxes in an integrated horizontal path covering about 80% of the basin. We present a preliminary analysis and characterization of the turbulent fluxes in the basin and we discuss the design of a multi-scale experimental and modeling approach to the study of ET that integrates point, spatial and volumetric in situ measurements, up to satellite scale observations. The ultimate focus of this exercise is to develop a consistent satellite-based ET retrieval approach.




## Imnavait Creek Topography, Experimental Layout & Instruments



The Imnavait Creek Basin is a small headwater basin of approximately 2 km<sup>2</sup>, located in the foothills of the Brooks Range (68°30'N, 149°15'W), 250 km south of the Arctic Ocean. The Imnavait Creek flows parallel to the Kuparuk River for 12 km before it joins the Kuparuk River that drains into the Arctic Ocean. The topographic elevation in this area ranges from 880 m at the outlet to 960 m (Trochim, 2009)

A preliminary experiment was carried out to characterize turbulent fluxes during the summer period from July to September 2009 including hydrological, meteorological and turbulent instruments.




Meteorological, Hydrological (left panel) and Turbulent Measurements in the Boundary Layer including sonic anemometers (middle panel) and scintillometer (right panel)


Upper panel (Kuparuk Basin, Alaskan North Slope)  
Lower panel (Imnavait Creek Basin)  
Experimental site and location of hydrological, meteorological and turbulent instruments)

## Turbulent fluxes punctual and horizontal path integrated

Surface Heat ( $\overline{w'\theta}$ ) and Momentum fluxes ( $\overline{u'w}$ ) derived by Eddy-Covariance from 3m – sonic anemometers tower-





Sensible heat flux derived from Boundary Layer Scintillometer 1 km path integration





## Satellite Remote Sensing Retrieval of Evapotranspiration- Two Source Energy Balance (TSEB) model

Symbol	Units	Description
$\rho_a$	kg/m <sup>3</sup>	Reflectivity of soil-canopy system
$\rho_s$	-	Priority-Taylor coefficient for canopy transpiration
$\alpha_s$	-	Soil heat flux coefficient
$\beta$	M/K	Slope of saturation vapor pressure vs. temperature curve
$\beta$	-	Leaf area index
$f_{veg}$	-	Fractional vegetation cover at view angle $\theta$
$f_{veg}$	-	Fraction of green vegetation
$Q$	W/m <sup>2</sup>	Soil heat flux
$\gamma$	M/K	Psychrometer constant
$H$	W/m <sup>2</sup>	System sensible heat flux
$H_c$	W/m <sup>2</sup>	Canopy sensible heat flux
$H_s$	W/m <sup>2</sup>	Soil sensible heat flux
$\lambda$	J/kg	Latent heat of vaporization
$\lambda E$	W/m <sup>2</sup>	System latent heat flux
$\lambda E_c$	W/m <sup>2</sup>	Canopy transpiration
$\lambda E_s$	W/m <sup>2</sup>	Soil transpiration
$\alpha$	deg	TIR sensor view angle from nadir
$R_a$	m <sup>2</sup>	Aerodynamic resistance for momentum
$R_c$	m <sup>2</sup>	Bulk two-sided leaf boundary layer resistance
$R_s$	m <sup>2</sup>	Soil boundary layer resistance
$R_{net}$	W/m <sup>2</sup>	Net radiation above canopy
$R_{net,c}$	W/m <sup>2</sup>	Net radiation divergence within canopy
$R_{net,s}$	W/m <sup>2</sup>	Net radiation above soil surface
$\rho_a$	J/K m <sup>3</sup>	Volumetric heat capacity of air
$T_a$	K	Aerodynamic temperature
$T_c$	K	Canopy temperature
$T_s$	K	Soil surface temperature
$T_{surf}(\theta)$	K	Surface radiometric temperature at view angle $\theta$

Preliminary results of recent efforts to model mass and energy fluxes in this arctic watershed using the USDA TSEB model (Anderson et al., 2008), which has been successfully applied at lower latitudes. The agreement between the modeled and observed fluxes may improve as we adjust the input vegetation parameters (leaf area index, green fraction, etc.) to values appropriate for the arctic, and as we identify arctic-specific modifications to the model that may better describe the energy flux in the very cold soils. Point-mode model results are used to tune the input parameters by comparison to the Arctic Observatory Network Imnavait Ridge station flux and met tower data (lower panel left), and these are then combined with satellite brightness temperature data to model the flux over an area that includes the station (lower panel right).




## Synoptic Flow Statistics & Meteorological Scales



Synoptic air mass in the Imnavait Basin is dominated by Southern flows with two secondary statistical modes WNW and ENE due to the basin orientation.

Morlet wavelet power spectrum demonstrate the occurrence of these two modes after the summer maxima when near-surface temperature approaches the freezing point.



## Summary

Local scale temporal and spatial variability of fluxes is induced by surface heterogeneity (water tracks), small scale local flows (downslope flows) and thermodynamic (late summer fog) processes in the Imnavait Basin.

Satellite retrieval reveals signatures according to basic thermodynamic processes in the basin.

This preliminary experiment demonstrates the need for a multi-scale flux observing system with volume control to fully capture the variability of fluxes. Single-point, horizontal path and volume-integrated measurements are necessary to account for space-time variability of evapotranspiration fluxes.

Consistent satellite retrieval rely on a realistic ground-truth calibration accounting for the physical properties of the basin.

## Acknowledgements

This research was supported by the Alaska NASA EPSCoR program award NNX07A037A, partially supported by the Department of Environmental Conservation of Alaska and the University Partnering for Operational support UPOS program under contract with the United States Air Force and the U.S. Army Research Laboratory and the Geophysical Institute of the University of Alaska Fairbanks.

## References

"Northern Research Basins Water Balance". D. Kane and D. Yang (Editors). Publication 290, 271 pages. International Association of Hydrological Sciences 2004.  
Trochim, E. (2009) "Modeling Discharge in Imnavait Basin, North Slope, Alaska". MS-Thesis University of Alaska Fairbanks.  
Anderson, M., Norman, J., Kustas, W., Houborg, R., Starks, P., and Agam, N. (2008) "A thermal-based remote sensing technique for routine mapping of land-surface carbon, water and energy fluxes from field to regional scales". Remote Sensing of Environment 112, 4227-4241.  
Brete-Harte, S., and Shaver, G. (2009) Climate and flux data from the International Study of Carbon and Energy Balances in Arctic Terrestrial Ecosystems project. University of Alaska Fairbanks, Institute of Arctic Biology, Arctic Observatory Network. URL: <http://aon.iab.uaf.edu/index.html>, variously paged. Data used with permission. Retrieved 9/23/2009.